Author index

Abubakar, Z. *see* Black, R.
Adams, A.N. *see* Kirby, M.J.
Aked, J. *see* Khan, S.H.
Aljanabi, S.M., Parmessur, Y., Moutia, Y., Saumtally, S. & Dookun, A. Further evidence of the association of a phytoplasma and a virus with yellow leaf syndrome in sugarcane, 628
Amorim, L. *see* Bassanezi, R.B.
Andrés, M.F., Romero, M.D., Montes, M.J. & Delibes, A. Genetic relationships and isozyme variability in the *Heterodera avenae* complex determined by isoelectrofocusing, 270
Anselmi, N. *see* Vettraino, A.M.
Anthony, F. *see* Bertrand, B.
Arraiano, L.S. *see* Brown, J.K.M.
Arraiano, L.S., Brading, P.A. & Brown, J.K.M. A detached seedling leaf technique to study resistance to *Mycosphaerella graminicola* (anamorph *Septoria tritici*) in wheat, 339
Arsenault, W.J. *see* Kimpinski, J.
see also Peters, R.D.
Arthofer, W. *see* Laimer Da Câmara Machado, M.
Ash, G.J. & Lanoiselet, V.M. First report of *Alternaria alternata* causing late blight of pistachio (*Pistacia vera*) in Australia, 803

Bakonyi, J., Nagy, Z.Á., Vajna, L. & Érsek, T. *Phytophthora nicotianae* causes blight of lily in Hungary, 795
Bandyopadhyay, R. *see* Dahlberg, J.A.
Barbara, D.J., Morton, A., Ramcharan, S., Cole, I.W., Phillips, A. & Knight, V.H. Occurrence and distribution of *Raspberry bushy dwarf virus* in commercial *Rubus* plantations in England and Wales, 747
Barbedette, S. *see* Ziadi, S.
Barbetti, M.J. *see* West, J.S.
Baron, F. *see* Manzanares-Dauleux, M.J.
Barreto, R.W. *see* Gonçalves, R.C.
Bassanezi, R.B., Amorim, L., Bergamin Filho, A., Hau, B. & Berger, R.D. Accounting for photosynthetic efficiency of bean leaves with rust, angular leaf spot and anthracnose to assess crop damage, 443
Bateman, G.L. *see* Dawson, W.A.J.M.
see also Dyer, P.S.; Turner, A.S.
Bates, J.A. *see* Taylor, E.J.A.
Bergamin Filho, A. *see* Bassanezi, R.B.
Berger, D.K. *see* Botha, W.J.
Berger, R.D. *see* Bassanezi, R.B.
Berrie, A.M. *see* Xu, X.-M.
Bertaccini, A. *see* Laimer Da Câmara Machado, M.
Bertelsen, J.R., De Neergaard, E. & Smedegaard-Petersen, V. Fungicidal effects of azoxystrobin and epoxiconazole on phyllosphere fungi, senescence and yield of winter wheat, 190
Bertrand, B., Anthony, F. & Lashermes, P. Breeding for resistance to *Meloidogyne exigua* in *Coffea arabica* by introgression of resistance genes of *Coffea canephora*, 637
Black, R., Seal, S., Abubakar, Z., Nono-Womdim, R. & Swai, I. Bacterial spot (*Xanthomonas campestris* pv. *vesicatoria*) of tomato and sweet pepper in Tanzania, 810
Blockley, A.L. & Mumford, R.A. The identification and isolation of *Impatiens necrotic spot virus* from prickly pear cactus (*Opuntia microdasys*), 805
Botha, W.J., Serfontein, S., Greyling, M.M. & Berger, D.K. Detection of *Xylophilus ampelinus* in grapevine cuttings using a nested polymerase chain reaction, 515
Bousset, L., Schaeffer, B. & De Vallavieille-Pope, C. Effect of early infections on pathotype frequencies in barley powdery mildew (*Blumeria graminis* f.sp. *hordei*) populations in field plots, 317
Brading, P.A. *see* Arraiano, L.S.
see also Brown, J.K.M.
Brasier, C.M. & Kirk, S.A. Comparative aggressiveness of standard and variant hybrid alder phytophthoras, *Phytophthora cambivora* and other *Phytophthora* species on bark of *Alnus*, *Quercus* and other woody hosts, 218
Breese, W.A. *see* Jones, E.S.
Broquaire, J.M. *see* Jarausch, W.
Brown, J.K.M. *see* Arraiano, L.S.
Brown, J.K.M., Kema, G.H.J., Forrer, H.-R., Verstappen, E.C.P., Arraiano, L.S., Brading, P.A., Foster, E.M., Fried, P.M. & Jenny, E. Resistance of wheat cultivars and breeding lines to septoria tritici blotch caused by isolates of *Mycosphaerella graminicola* in field trials, 325
Brun, H., Ruer, D., Levivier, S., Somda, I., Renard, M. & Chèvre, A.M. Presence in *Leptosphaeria maculans* populations of isolates virulent on resistance introgressed into *Brassica napus* from the *B. nigra* B genome, 69
Buonaurio, R., Stravato, V.M. & Cappelli, C. Brown spot caused by *Sphingomonas* sp. on yellow Spanish melon fruits in Spain, 397
Burgess, T. *see* Maseko, B.
Burr, H.N. *see* Wong, F.P.
Buttgereitt, A. *see* Ravi, K.S.

Cappelli, C. *see* Buonaurio, R.
Capron, G. *see* Robin, C.
Carimi, F. *see* D'Onghia, A.M.
Cartia, G. *see* Di Primo, P.
Castilla, N.P. *see* Savary, S.
Castro, L.H.R. *see* Tessmann, D.J.
Chacón, G. *see* Trognitz, B.R.
Chèvre, A.M. *see* Brun, H.
Cho, H.S. *see* Yu, S.H.
Christian, D.G. *see* Etheridge, J.V.
Christiansen, E. *see* Krokene, P.
Claussen, W. *see* Grote, D.
Clewer, A. *see* Sijaona, M.E.R.
Clover, G.R.G., Ratti, C. & Henry, C.M. Molecular characterization and detection of European isolates of *Soil-borne wheat mosaic virus*, 761
Coghe, F. *see* Roggero, P.
Cole, I.W. *see* Barbara, D.J.
Collinge, D.B. *see* Haugaard, H.
Colquhoun, I.J. *see* Wilkinson, C.J.
Colvin, J. *see* Zhang, X.-S.
Cook, R.T.A. First report in England of changes in the susceptibility of *Puccinia horiana*, the cause of chrysanthemum white rust, to triazole and strobilurin fungicides, 792
Cotton, S. *see* Monger, W.A.
Coutinho, T. *see* Maseko, B.
Coutinho, T.A. *see* Slippers, B.
see also Venter, M.

D'Onghia, A.M., Carimi, F., De Pasquale, F., Djelouah, K. & Martelli, G.P. Elimination of *Citrus psorosis virus* by somatic embryogenesis from stigma and style cultures, 266
Dahlberg, J.A., Bandyopadhyay, R., Rooney, W.L., Odvody, G.N. & Madera-Torres, P. Evaluation of sorghum germplasm used in US breeding programmes for sources of sugary disease resistance, 681
Danet, J.L. *see* Jarausch, W.
Davey, L. *see* Etheridge, J.V.
Dawson, W.A.J.M. & Bateman, G.L. Fungal communities on roots of wheat and barley and effects of seed treatments containing fluquinconazole applied to control take-all, 75

© 2001 BSPP

De Lacy Costello, B.P.J., Evans, P., Ewen, R.J., Gunson, H.E., Jones, P.R.H., Ratcliffe, N.M. & Spencer-Phillips, P.T.N. Gas chromatography–mass spectrometry analyses of volatile organic compounds from potato tubers inoculated with *Phytophthora infestans* or *Fusarium coeruleum*, 489
De Neergaard, E. *see* Bertelsen, J.R.
De Pasquale, F. *see* D'Onghia, A.M.
De Vallavieille-Pope, C. *see* Bousset, L.
Delibes, A. *see* Andrés, M.F.
Dell, B. *see* McKenna, F.
see also Wilkinson, C.J.
Derks, J.H.J. *see* Janse, J.D.
Deshmukh, S. *see* Ravi, K.S.
Desprez-Loustau, M.L. *see* Robin, C.
Devonshire, B.J. *see* Pandit, M.K.
Di Primo, P., Cartia, G. & Katan, T. Vegetative compatibility and heterokaryon stability in *Fusarium oxysporum* f.sp. *radicis-lycopersici* from Italy, 371
Dianese, J.C. *see* Tessmann, D.J.
Dickinson, M.J. *see* Steele, K.A.
Divaret, I. *see* Manzanares-Dauleux, M.J.
Djelouah, K. *see* D'Onghia, A.M.
Dookun, A. *see* Aljanabi, S.M.
Dosba, F. *see* Jarausch, W.
Drenth, A. *see* Linde, C.
Dyer, P.S., Bateman, G.L. & Wood, H.M. Development of apothecia of the eyespot pathogen *Tapesia* on cereal crop stubble residue in England, 356

Edwards, S.G. *see* Turner, A.S.
Egan, D. *see* Qu, X.-S.
El-Muadhidi, M.A. *see* Makkouk, K.M.
El-Tarabily, K.A. *see* McKenna, F.
Ena, S. *see* Roggero, P.
Erper, I. *see* Karaca, G.H.
Errampalli, D., Saunders, J.M. & Holley, J.D. Emergence of silver scurf (*Helminthosporium solani*) as an economically important disease of potato, 141
Érsek, T. *see* Bakonyi, J.
Etheridge, J.V., Davey, L. & Christian, D.G. First report of *Rhizoctonia cerealis* causing sharp eyespot in *Panicum virgatum* in the UK, 807
Evans, N. *see* Gilles, T.
Evans, P. *see* De Lacy Costello, B.P.J.
Ewen, R.J. *see* De Lacy Costello, B.P.J.

Farreyrol, K., Pearson, M.N., Grisoni, M. & Leclercq-Le Quillec, F. Severe stunting of *Vanilla tahitensis* in French Polynesia caused by *Cucumber mosaic virus* (CMV), and the detection of the virus in *V. fragrans* in Reunion Island, 414
Fitch, M.M.M., Lehrer, A.T., Komor, E. & Moore, P.H. Elimination of *Sugarcane yellow leaf virus* from infected sugarcane plants by meristem tip culture visualized by tissue blot immunoassay, 676
Fitt, B.D.L. *see also* Gilles, T.; Papastamati, K.; Toscano-Underwood, C.; West, J.S.
Flier, W.G., Turkensteen, L.J., Van den Bosch, G.B.M., Vereijken, P.F.G. & Mulder, A. Differential interaction of *Phytophthora infestans* on tubers of potato cultivars with different levels of blight resistance, 292
Forrer, H.-R. *see* Brown, J.K.M.
Foster, E.M. *see* Brown, J.K.M.
Foster, G.D. *see* Monger, W.A.
Fránová, J. *see* Přibylová, J.
Fried, P.M. *see* Brown, J.K.M.
Froidefond, G. *see* Guerin, L.

Garnier, M. *see* Jarausch, W.
Gefen, T. *see* Hadas, R.
Georgiou, C.D. & Petropoulou, K.P. Effect of the antioxidant ascorbic acid on sclerotial differentiation in *Rhizoctonia solani*, 594
Giesbrecht, F.G. *see* Whitaker, T.B.
Gilbert, R.L. *see* Nikandrow, A.
Gilles, T., Fitt, B.D.L., Welham, S.J., Evans, N., Steed, J.M. & Jeger, M.J. Modelling the effects of temperature and wetness duration on development of light leaf spot on oilseed rape leaves inoculated with *Pyrenopeziza brassicae* conidia, 42
Gladders, P. *see* Papastamati, K.
Godard, J.F. *see* Ziadi, S.
Gogoi, R., Singh, D.V. & Srivastava, K.D. Phenols as a biochemical basis of resistance in wheat against Karnal bunt, 470
Gonçalves, R.C. & Barreto, R.W. Root and stem rot of *Schefflera arboricola* in Brazil, caused by *Fusarium solani*, 794
Gorkink, R.F.J. *see* Janse, J.D.
Greyling, M.M. *see* Botha, W.J.
Grisoni, M. *see* Farreyrol, K.
Grote, D. & Çlaussen, W. Severity of root rot on tomato plants caused by *Phytophthora nicotianae* under nutrient- and light-stress conditions, 702
Guerin, L. *see* Xu, X.-M.
Guerin, L., Froidefond, G. & Xu, X.-M. Seasonal patterns of dispersal of ascospores of *Cryphonectria parasitica* (chestnut blight), 717
Gunning, D.A. *see* Nikandrow, A.
Gunson, H.E. *see* De Lacy Costello, B.P.J.

Hadas, R., Kritzman, G., Gefen, T. & Manulis, S. Detection, quantification and characterization of *Erwinia carotovora* ssp. *carotovora* contaminating pepper seeds, 117
Hamaide, N. *see* Munaut, F.
Hanzer, V. *see* Laimer Da Câmara Machado, M.
Hardwick, N.V. Fifty years of *Plant Pathology*, 1
Hardwick, N.V., Jones, D.R. & Slough, J.E. Factors affecting diseases of winter wheat in England and Wales, 453
Hardwick, N.V., Jones, D.R. & Slough, J.E. Erratum: Factors affecting diseases of winter wheat in England and Wales, 1989–98, 650
Hardy, G.E.St J. *see* McKenna, F.
see also Wilkinson, C.J.
Harris, D.C. *see* Xu, X.-M.
Hau, B. *see* Bassanezi, R.B.
Haugaard, H., Lyngs Jørgensen, H.J., Lyngkjær, M.F., Smedegaard-Petersen, V. & Collinge, D.B. Control of *Blumeria graminis* f.sp. *hordei* by treatment with mycelial extracts from cultured fungi, 552
Hazanovsky, M. *see* Tsror (Lahkim), L.
Henry, C.M. *see* Clover, G.R.G.
Hick, A. *see* Monda, E.O.
Holley, J.D. *see* Errampalli, D.
Holmes, J.M. *see* Wilkinson, C.J.
Holt, J. *see* Zhang, X.-S.
Hovmøller, M.S. Disease severity and pathotype dynamics of *Puccinia striiformis* f.sp. *tritici* in Denmark, 181
Huang, R. & Tu, J.C. Effects of nutrient solution pH on the survival and transmission of *Clavibacter michiganensis* ssp. *michiganensis* in hydroponically grown tomatoes, 503
Hullé, M. *see* Wicker, E.
Humphreys, E. *see* Steele, K.A.
Husted, K. *see* Nicolaisen, M.

Isaac, A.M. *see* Monger, W.A.

Jackson, T.J. *see* Wilkinson, C.J.
Janse van Rensburg, J.C., Labuschagne, N. & Nemec, S. Occurrence of *Fusarium*-produced naphthazarins in citrus trees and sensitivity of rootstocks to isomarticin in relation to citrus blight, 258
Janse, J.D., Rossi, M.P., Gorkink, R.F.J., Derks, J.H.J., Swings, J., Janssens, D. & Scortichini, M. Bacterial leaf blight of strawberry *(Fragaria (x) ananassa)* caused by a pathovar of *Xanthomonas arboricola*, not similar to *Xanthomonas fragariae* Kennedy & King. Description of the causal organism as *Xanthomonas arboricola* pv. *fragariae* (pv. nov., comb. nov.), 653
Janssens, D. *see* Janse, J.D.
Jarausch, W., Danet, J.L., Labonne, G., Dosba, F., Broquaire, J.M., Saillard, C. & Garnier, M. Mapping the spread of apricot chlorotic leaf roll (ACLR) in southern France and implication of *Cacopsylla pruni* as a vector of European stone fruit yellows (ESFY) phytoplasmas, 782

Jedryczka, M. *see* Toscano-Underwood, C.
Jeger, M.J. *see* Gilles, T.
Jenny, E. *see* Brown, J.K.M.
Jiménez-Díaz, R.M. *see* Mercado-Blanco, J.
Johansson, A.S. *see* Whitaker, T.B.
Jones, D.R. *see* Hardwick, N.V.
Jones, E.S., Breese, W.A. & Shaw, D.S. Inoculation of pearl millet with the downy mildew pathogen, *Sclerospora graminicola*: chilling inoculum to delay zoospore release and avoid spray damage to zoospores, 310
Jones, P. *see* Pandit, M.K.
Jones, P.R.H. *see* De Lacy Costello, B.P.J.

Karaca, G.H. & Erper, I. First report of *Pestalotiopsis guepinii* causing twig blight on hazelnut and walnut in Turkey, 415
Katan, T. *see* Di Primo, P.
Kavanagh, J.A. *see* Qu, X.-S.
Kema, G.H.J. *see* Brown, J.K.M.
Kenyon, D.M. *see* Taylor, E.J.A.
Khan, S.H., Aked, J. & Magan, N. Control of the anthracnose pathogen of banana (*Colletotrichum musae*) using antioxidants alone and in combination with thiabendazole or imazalil, 601
Kharbanda, P.D. *see* West, J.S.
Kimpinski, J., Arsenault, W.J. & Sturz, A.V. Differential effect of nematicide treatments on tuber yields in early- and late-maturing potato cultivars, 509
Kirby, M.J. *see* Kirby, M.J.
Kirby, M.J., Kirby, M.J. & Adams, A.N. Occurrence of *Cherry virus A* in the UK, 801
Kirk, S.A. *see* Brasier, C.M.
Kiss, L. & Szentiványi, O. Infection of bean with cucumber powdery mildew, *Podosphaera fusca*, 411
Kitkaru, A.S. *see* Ravi, K.S.
Knight, V.H. *see* Barbara, D.J.
Kolmer, J.A. & Liu, J.Q. Simple inheritance of partial resistance to leaf rust in two wheat cultivars, 546
Kolnaar, R.W. & Van den Bosch, F. Effect of temperature on epidemiological parameters of *Puccinia lagenophorae*, 363
Komor, E. *see* Fitch, M.M.M.
Kostova, D., Masenga, V., Milne, R.G. & Lisa, V. First report of *Eggplant mottled dwarf virus* in cucumber and pepper in Bulgaria, 804
Kritzman, G. *see* Hadas, R.
Krokene, P. & Solheim, H. Loss of pathogenicity in the blue-stain fungus *Ceratocystis polonica*, 497
Krokene, P. *see* Solheim, H.
Krokene, P., Solheim, H. & Christiansen, E. Induction of disease resistance in Norway spruce (*Picea abies*) by necrotizing fungi, 230
Kumari, S.G. *see* Makkouk, K.M.
Kurt, S. & Soylu, S. First report of powdery mildew on mulberry caused by *Phyllactinia guttata* in the eastern Mediterranean region of Turkey, 797

Labonne, G. *see* Jarausch, W.
Labuschagne, N. *see* Janse van Rensburg, J.C.
Lahert, H. *see* Qu, X.-S.
Laimer Da Câmara Machado, M., Paltrinieri, S., Hanzer, V., Arthofer, W., Strommer, S., Martini, M., Pondrelli, M. & Bertaccini, A. Presence of European stone fruit yellows (ESFY or 16SrX-B) phytoplasmas in apricots in Austria, 130
Långström, B. *see* Solheim, H.
Lanoiselet, V.M. *see* Ash, G.J.
Lashermes, P. *see* Bertrand, B.
Lawler, M.A.C.B. *see* Nikandrow, A.
Le Corre, D. *see* Ziadi, S.
Leclercq-Le Quillec, F. *see* Farreyrol, K.
Lee, D. *see* Taylor, E.J.A.
Lee, H.B. *see* Yu, S.H.
Legg, J.P., Okao-Okuja, G., Mayala, R. & Muhinyuza, J.-B. Spread into Rwanda of the severe cassava mosaic virus disease pandemic and associated Uganda variant of *East African cassava mosaic virus* (EACMV-Ug), 796
Lehrer, A.T. *see* Fitch, M.M.M.
Lenzi, R. *see* Roggero, P.
Lesemann, D.E. *see* Ravi, K.S.
Levivier, S. *see* Brun, H.
Lindbeck, K.D. *see* Nikandrow, A.
Linde, C., Soo, S.H. & Drenth, A. Sexual recombination in *Phytophthora cinnamomi in vitro* and aggressiveness of single-oospore progeny to *Eucalyptus*, 97
Lisa, V. *see* Kostova, D.
Liu, J.Q. *see* Kolmer, J.A.
Lonsdale, D. *see* Zoina, A.
Lyngkjær, M.F. *see* Haugaard, H.
Lyngs Jørgensen, H.J. *see* Haugaard, H.

Maddison, A. *see* Sijaona, M.E.R.
Madera-Torres, P. *see* Dahlberg, J.A.
Magan, N. *see* Khan, S.H.
Makkouk, K.M., El-Muadhidi, M.A. & Kumari, S.G. First record of *Beet western yellows virus*, *Chickpea chlorotic dwarf virus* and *Faba bean necrotic yellows virus* affecting faba bean (*Vicia faba*) crops in Iraq, 793
Makkouk, K.M., Najar, A. & Kumari, S.G. The first record of *Barley yellow dwarf virus* and *Cereal yellow dwarf virus* in Tunisia, 806
Malone, A. *see* Peters, R.D.
Mansfield, J.W. *see* Sijaona, M.E.R.
Manulis, S. *see* Hadas, R.
Manzanares-Dauleux, M.J., Divaret, I., Baron, F. & Thomas, G. Assessment of biological and molecular variability between and within field isolates of *Plasmodiophora brassicae*, 165
Maraite, H. *see* Munaut, F.
Marshall, J. *see* Turner, A.S.
Martelli, G.P. *see* D'Onghia, A.M.
Martini, M. *see* Laimer Da Câmara Machado, M.
Maseko, B., Burgess, T., Coutinho, T. & Wingfield, M. First report of *Phytophthora nicotianae* associated with *Eucalyptus* die-back in South Africa, 413
Masenga, V. *see* Kostova, D.
see also Roggero, P.
Matheson, B.G. *see* Peters, R.D.
Mathur, S.B. *see* Tagne, A.
Mayala, R. *see* Legg, J.P.
McComb, J.A. *see* Wilkinson, C.J.
McKenna, F., El-Tarabily, K.A., Hardy, G.E.St J. & Dell, B. Novel *in vivo* use of a polyvalent *Streptomyces* phage to disinfest *Streptomyces scabies*-infected seed potatoes, 666
Mercado-Blanco, J., Rodríguez-Jurado, D., Pérez-Artés, E. & Jiménez-Díaz, R.M. Detection of the nondefoliating pathotype of *Verticillium dahliae* in infected olive plants by nested PCR, 609
Metcalf, D.A. & Wilson, C.R. The process of antagonism of *Sclerotium cepivorum* in white rot affected onion roots by *Trichoderma koningii*, 249
Milne, R.G. *see* Kostova, D.
Miranda, A.C. *see* Tessmann, D.J.
Monda, E.O., Sanders, F.E. & Hick, A. Infection and colonization of bean leaf by *Phaeoisariopsis griseola*, 103
Monger, W.A., Seal, S., Cotton, S. & Foster, G.D. Identification of different isolates of *Cassava brown streak virus* and development of a diagnostic test, 768
Monger, W.A., Seal, S., Isaac, A.M. & Foster, G.D. Molecular characterization of the *Cassava brown streak virus* coat protein, 527
Monot, C. *see* Ziadi, S.
Montes, M.J. *see* Andrés, M.F.
Moore, P.H. *see* Fitch, M.M.M.
Mordechi-Lebiush, S. *see* Tsror (Lahkim), L.
Morgan, L.W. *see* Turner, A.S.
Morreale, G. *see* Taylor, E.J.A.
Morton, A. *see* Barbara, D.J.
Moutia, Y. *see* Aljanabi, S.M.
Muhinyuza, J.-B. *see* Legg, J.P.
Mukhopadhyay, S. *see* Pandit, M.K.
Mulder, A. *see* Flier, W.G.
Mumford, R.A. *see* Blockley, A.L.
Munaut, F., Hamaide, N. & Maraite, H. Molecular and morphological characterization of *Colletotrichum gloeosporioides* from native Mexican *Stylosanthes* species,

383
Murray, G. *see* Sosnowski, M.
Murray, G.M. *see* Nikandrow, A.

Nagy, Z.Á. *see* Bakonyi, J.
Najar, A. *see* Makkouk, K.M.
Nath, P.S. *see* Pandit, M.K.
Natili, G. *see* Vettraino, A.M.
Nemec, S. *see* Janse van Rensburg, J.C.
Nicholson, P. *see* Turner, A.S.
Nicolaisen, M., Rasmussen, H.N., Husted, K. & Nielsen, S.L. Reverse transcription–detection of immobilized, amplified product in a one-phase system (RT-DIAPOPS) for the detection of potato virus Y, 124
Nielsen, S.L. *see* Nicolaisen, M.
Nikandrow, A., Gilbert, R.L., Gunning, D.A., Lawler, M.A.C.B., Lindbeck, K.D. & Murray, G.M. First report of *Phytophthora* sp. causing a root and basal stem rot of narrowleaf lupin in Australia, 811
Nono-Womdim, R. *see* Black, R.
Nuttall, M. *see* Turner, A.S.

Odvody, G.N. *see* Dahlberg, J.A.
Okao-Okuja, G. *see* Legg, J.P.
Orrillo, M. *see* Trognitz, B.R.

Paltrinieri, S. *see* Laimer Da Câmara Machado, M.
Pandit, M.K., Nath, P.S., Mukhopadhyay, S., Devonshire, B.J. & Jones, P. First report of *Dasheen mosaic virus* in elephant foot yam in India, 802
Papastamati, K., Welham, S.J., Fitt, B.D.L. & Gladders, P. Modelling the progress of light leaf spot (*Pyrenopeziza brassicae*) on winter oilseed rape (*Brassica napus*) in relation to leaf wetness and temperature, 154
Parmessur, Y. *see* Aljanabi, S.M.
Parry, D.W. *see* Turner, A.S.
Pearson, M.N. *see* Farreyrol, K.
Peluso, R. *see* Zoina, A.
Perez, S. *see* Trognitz, B.R.
Pérez-Artés, E. *see* Mercado-Blanco, J.
Peters, R.D., Sturz, A.V., Matheson, B.G., Arsenault, W.J. & Malone, A. Metalaxyl sensitivity of isolates of *Phytophthora erythroseptica* in Prince Edward Island, 302
Peterson, G.L. *see* Whitaker, T.B.
Petrzik, K. *see* Přibylová, J.
Petropoulou, K.P. *see* Georgiou, C.D.
Phillips, A. *see* Barbara, D.J.
Pondrelli, M. *see* Laimer Da Câmara Machado, M.
Portal, L. *see* Trognitz, B.R.
Přibylová, J., Špak, J., Fránová, J. & Petrzik, K. Association of aster yellows subgroup 16SrI-B phytoplasmas with a disease of *Rehmannia glutinosa* var. *purpurea*, 776

Qu, X.-S., Kavanagh, J.A., Egan, D. & Lahert, H. Studies on amoebae and cysts associated with the isolation of *Spongospora subterranea* f.sp. *subterranea in vitro*, 420

Raio, A. *see* Zoina, A.
Ramcharan, S. *see* Barbara, D.J.
Ramón, P. *see* Trognitz, B.R.
Ramsey, M. *see* Sosnowski, M.
Rasmussen, H.N. *see* Nicolaisen, M.
Ratcliffe, N.M. *see* De Lacy Costello, B.P.J.
Ratti, C. *see* Clover, G.R.G.
Ravi, K.S., Buttgereitt, A., Kitkaru, A.S., Deshmukh, S., Lesemann, D.E. & Winter, S. Sunflower necrosis disease from India is caused by an ilarvirus related to *Tobacco streak virus*, 800
Renard, M. *see* Brun, H.
Robin, C., Capron, G. & Desprez-Loustau, M.L. Root infection by *Phytophthora cinnamomi* in seedlings of three oak species, 708
Robinson, J.D. *see* Xu, X.-M.
Rodgers-Gray, B.S. & Shaw, M.W. The effect of incorporating straw or manure into the soil on the natural microflora of winter wheat, 537
Rodríguez-Jurado, D. *see* Mercado-Blanco, J.
Roggero, P., Masenga, V., Lenzi, R., Coghe, F., Ena, S. & Winter, S. First report of *Pepino mosaic virus* in tomato in Italy, 798
Román, C. *see* Trognitz, B.R.
Romero, M.D. *see* Andrés, M.F.
Rooney, W.L. *see* Dahlberg, J.A.
Rossi, M.P. *see* Janse, J.D.
Rouxel, F. *see* Wicker, E.
Ruer, D. *see* Brun, H.

Şahin, F. Pepper races 7, 8, and 10 of *Xanthomonas axonopodis* pv. *vesicatoria* isolated from diseased pepper plants in Turkey, 809
Şahin, F. Severe outbreak of bacterial speck, caused by *Pseudomonas syringae* pv. *tomato* on field-grown tomatoes in eastern Anatolia region of Turkey, 799
Saillard, C. *see* Jarausch, W.
Sanders, F.E. *see* Monda, E.O.
Saumtally, S. *see* Aljanabi, S.M.
Saunders, J.M. *see* Errampalli, D.
Savary, S., Castilla, N.P. & Willocquet, L. Analysis of the spatio-temporal structure of rice sheath blight epidemics in a farmer's field, 53
Schaeffer, B. *see* Bousset, L.
Schiff, C.L., Wilson, I.W. & Somerville, S.C. Polygenic powdery mildew disease resistance in *Arabidopsis thaliana*: quantitative trait analysis of the accession Warschau-1, 690
Scortichini, M. *see* Janse, J.D.
Scott, E. *see* Sosnowski, M.
Seal, S. *see* Black, R.
see also Monger, W.A.
Serfontein, S. *see* Botha, W.J.
Shattock, Richard C. *see* Zoina, A.
Shaw, D.S. *see* Jones, E.S.
Shaw, M.W. *see* Rodgers-Gray, B.S.
Shearer, B.L. *see* Wilkinson, C.J.
Sijaona, M.E.R. & Mansfield, J.W. Variation in the response of cashew genotypes to the targeted application of fungicide to flower panicles for control of powdery mildew disease, 244
Sijaona, M.E.R., Clewer, A., Maddison, A. & Mansfield, J.W. Comparative analysis of powdery mildew development on leaves, seedlings and flower panicles of different genotypes of cashew, 234
Silué, D. *see* Ziadi, S.
Singh, D.V. *see* Gogoi, R.
Sivan, S. *see* Tsror (Lahkim), L.
Slippers, B., Wingfield, M.J., Coutinho, T.A. & Wingfield, B.D. Population structure and possible origin of *Amylostereum areolatum* in South Africa, 206
Slough, J.E. *see* Hardwick, N.V.
Smedegaard-Petersen, V. *see* Bertelsen, J.R.
see also Haugaard, H.
Solheim, H. *see* Krokene, P.
Solheim, H., Krokene, P. & Långström, B. Effects of growth and virulence of associated blue-stain fungi on host colonization behaviour of the pine shoot beetles *Tomicus minor* and *T. piniperda*, 111
Somda, I. *see* Brun, H.
Somerville, S.C. *see* Schiff, C.L.
Soo, S.H. *see* Linde, C.
Sosnowski, M., Ramsey, M., Murray, G., Scott, E. & Wilmshurst, C. Symptoms of blackleg (*Leptosphaeria maculans*) on the roots of canola in Australia, 808
Soylu, S. *see* Kurt, S.
Špak, J. *see* Přibylová, J.
Spasiano, A. *see* Zoina, A.
Spencer-Phillips, P.T.N. *see* De Lacy Costello, B.P.J.
Srivastava, K.D. *see* Gogoi, R.
Steed, J.M. *see* Gilles, T.
Steele, K.A., Humphreys, E., Wellings, C.R. & Dickinson, M.J. Support for a stepwise mutation model for pathogen evolution in Australasian *Puccinia striiformis* f.sp. *tritici* by use of molecular markers, 174
Stevens, E.A. *see* Taylor, E.J.A.

Stravato, V.M. *see* Buonaurio, R.
Strommer, S. *see* Laimer Da Câmara Machado, M.
Sturz, A.V. *see* Kimpinski, J.
see also Peters, R.D.
Swai, I. *see* Black, R.
Swings, J. *see* Janse, J.D.
Szentiványi, O. *see* Kiss, L.

Tagne, A. & Mathur, S.B. First report of chlorotic spot of maize caused by *Pestalotiopsis neglecta*, 791
Taylor, E.J.A., Stevens, E.A., Bates, J.A., Morreale, G., Lee, D., Kenyon, D.M. & Thomas, J.E. Rapid-cycle PCR detection of *Pyrenophora graminea* from barley seed, 347
Tessmann, D.J., Dianese, J.C., Miranda, A.C. & Castro, L.H.R. Epidemiology of a Neotropical rust *(Puccinia psidii)*: periodical analysis of the temporal progress in a perennial host *(Syzygium jambos)*, 725
Thomas, G. *see* Manzanares-Dauleux, M.J.
Thomas, J.E. *see* Taylor, E.J.A.
Todd, A.D. *see* Toscano-Underwood, C.
see also Turner, A.S.
Toscano-Underwood, C., West, J.S., Fitt, B.D.L., Todd, A.D. & Jedryczka, M. Development of phoma lesions on oilseed rape leaves inoculated with ascospores of A-group or B-group *Leptosphaeria maculans* (stem canker) at different temperatures and wetness durations, 28
Trognitz, B.R., Orrillo, M., Portal, L., Román, C., Ramón, P., Perez, S. & Chacón, G. Evaluation and analysis of reduction of late blight disease in a diploid potato progeny, 281
Tsror (Lahkim), L. & Hazanovsky, M. Effect of coinoculation by *Verticillium dahlie* and *Colletotrichum coccodes* on disease symptoms and fungal colonization in four potato cultivars, 483
Tsror (Lahkim), L., Hazanovsky, M., Mordechi-Lebiush, S. & Sivan, S. Aggressiveness of *Verticillium dahliae* isolates from different vegetative compatibility groups to potato and tomato, 477
Tu, J.C. *see* Huang, R.
Turkensteen, L.J. *see* Flier, W.G.
Turner, A.S., Nicholson, P., Edwards, S.G., Bateman, G.L., Morgan, L.W., Todd, A.D., Parry, D.W., Marshall, J. & Nuttall, M. Evaluation of diagnostic and quantitative PCR for the identification and severity assessment of eyespot and sharp eyespot in winter wheat, 463
Tynan, K.M. *see* Wilkinson, C.J.

Vajna, L. *see* Bakonyi, J.
Van den Bosch, F. *see* Kolnaar, R.W.
Van den Bosch, G.B.M. *see* Flier, W.G.
Vannini, A. *see* Vettraino, A.M.
Venter, M., Wingfield, M.J., Coutinho, T.A. & Wingfield, B.D. Molecular characterization of *Endothia gyrosa* isolates from *Eucalyptus* in South Africa and Australia, 211
Vereijken, P.F.G. *see* Flier, W.G.
Verstappen, E.C.P. *see* Brown, J.K.M.
Vettraino, A.M., Natili, G., Anselmi, N. & Vannini, A. Recovery and pathogenicity of *Phytophthora* species associated with a resurgence of ink disease in *Castanea sativa* in Italy, 90

Welham, S.J. *see* Gilles, T.
see also Papastamati, K.
Wellings, C.R. *see* Steele, K.A.
West, J.S. *see* Toscano-Underwood, C.
West, J.S., Kharbanda, P.D., Barbetti, M.J. & Fitt, B.D.L. Epidemiology and management of *Leptosphaeria maculans* (phoma stem canker) on oilseed rape in Australia, Canada and Europe, 10
Whitaker, T.B., Wu, J., Peterson, G.L., Giesbrecht, F.G. & Johansson, A.S. Variability associated with the official USDA sampling plan used to inspect export wheat shipments for *Tilletia controversa* spores, 755
Wicker, E., Hullé, M. & Rouxel, F. Pathogenic characteristics of isolates of *Aphanomyces euteiches* from pea in France, 433
Wilcox, W.F. *see* Wong, F.P.
Wilkinson, C.J., Holmes, J.M., Dell, B., Tynan, K.M., McComb, J.A., Shearer, B.L., Colquhoun, I.J. & Hardy, G.E.St J. Effect of phosphite on *in planta* zoospore production of *Phytophthora cinnamomi*, 587
Wilkinson, C.J., Shearer, B.L., Jackson, T.J. & Hardy, G.E.St J. Variation in sensitivity of Western Australian isolates of *Phytophthora cinnamomi* to phosphite *in vitro*, 83
Willocquet, L. *see* Savary, S.
Wilmshurst, C. *see* Sosnowski, M.
Wilson, C.R. Resistance to infection and translocation of *Tomato spotted wilt virus* in potatoes, 402
Wilson, C.R. *see* Metcalf, D.A.
Wilson, I.W. *see* Schiff, C.L.
Wingfield, B.D. *see* Slippers, B.
see also Venter, M.
Wingfield, M. *see* Maseko, B.
Wingfield, M.J. *see* Slippers, B.
see also Venter, M.
Winter, S. *see* Ravi, K.S.
see also Roggero, P.
Wong, F.P., Burr, H.N. & Wilcox, W.F. Heterothallism in *Plasmopara viticola*, 427
Wood, H.M. *see* Dyer, P.S.
Wu, J. *see* Whitaker, T.B.

Xu, X.-M *see* Guerin, L.
Xu, X.-M., Guerin, L. & Robinson, J.D. Effects of temperature and relative humidity on conidial germination and viability, colonization and sporulation of *Monilinia fructigena*, 561
Xu, X.-M., Robinson, J.D., Berrie, A.M. & Harris, D.C. Spatio-temporal dynamics of brown rot (*Monilinia fructigena*) on apple and pear, 569

Yu, S.H., Cho, H.S. & Lee, H.B. First report of alternaria leaf blight of blackberry-lily caused by *Alternaria iridicola* in Korea, 412

Zhang, X.-S., Holt, J. & Colvin, J. Synergism between plant viruses: a mathematical analysis of the epidemiological implications, 732
Ziadi, S., Barbedette, S., Godard, J.F., Monot, C., Le Corre, D. & Silué, D. Production of pathogenesis-related proteins in the cauliflower (*Brassica oleracea* var. *botrytis*)-downy mildew (*Peronospora parasitica*) pathosystem treated with acibenzolar-S-methyl, 579
Zoina, A., Lonsdale, D. & Shattock, R.C. Laboratory guide to the identification of plant pathogenic bacteria, third edition. 2. *Ganoderma* diseases of perannial crops. 3. Stem rust wheat: from ancient enemy to modern foe., 812
Zoina, A., Raio, A., Peluso, R. & Spasiano, A. Characterization of agrobacteria from weeping fig (*Ficus benjamina*), 620

Plant Pathology (2001) **50**, 820–822

Keyword index

16S rDNA sequence analysis, 397
16S rDNA sequence, 776
3 glucanase, 579

A group and B group, 10
A1/A2 mating tape, 83
adult plant resistance, 546
AFLP, 174
AFLPs, 637
aggregation, 569
aggressiveness, 97, 218, 292
Aggrobacterium spp., 620
agronomy, 537
Alnus, 218
amoebae, 420
Amylostereum, 206
Anacardium occidentale, 234
angular leaf spot, 103
antioxidant, 593
apothecia, 356
apple, 561, 569
apricot decline and dieback, 130
apricot, 782
ARIMA model, 717
ascorbic acid, 593
ascospores, 28, 154, 717
aseptic plantlets, 483
asexual generations, 42
Auchenorryhncha, 782
azoxystrobin, 190

bacterial blight, 515
bait-plant trials, 782
banana, 601
Banksia, 587
bark lesions, 218
barley, 75, 552
benzothiadiazole, 579
beta-1, 579
biological control, 552, 666
biopesticide, 552
bioreactor, 666
biotic and abiotic stress, 702
black dot, 483
blackleg, 10, 69
blue-stain fungi, 111
blue-stain fungus, 497
Blumeria graminis, 317, 552
breeding, 637
broad-sense heritability, 281
cambivora, 90

canola, 10
Capiscum annuum, 117
cashew, 23, 244
Cassava brown steak virus, 527, 768
cassava mosaic virus disease, 732
cassava, 527
Castanea sativa, 90
Castanea, 218
cellular fatty acids, 653
Ceratocystis polonica, 230, 497
cereal cyst nematode, 270
cereal diseases, 453
certification, 402
characterization, 761
chemotaxonomy, 653
Chestnut blight, 717
chitinase, 249, 579
Citrus psorosis virus, 266
citrus, 258
citrus, 266
Claviceps africana, 681
clubroot, 165
coexistence, 732
Coffea, 637
Colletotrichum lindemuthianum, 443
Colletotrichum, 601
common scab, 666
competitive exclusion, 732
conidia, 154, 601
crown canker, 10
crown gall, 620
crucifers, 165
Cryphonectria cubensis, 211
Cryphonectria parasitica, 211, 717
Cucumis melo, var. *Inodorus*, 397
cultivar-by-isolate interactions, 325, 339
cultivars, 453
cysts, 420

DAC-ELISA, 702
detached leaf technique, 339
detection, 402, 761
diagnostic assays, 463
diagnostics, 515
DIAPOPS, 124
differentiation, 593
disease forecasting, 42
disease incidence, 53, 117, 181
disease management, 141
disease progress, 42, 154
disease resistance, 690
disease severity, 181
disease thresholds, 463
dissemination, 717
DNA:DNA hybridisation, 653
downy mildew, 310, 427
dwarf bunt, 755

early- and late- maturing potato cultivars, 509
early dying of potato, 477
electron microscopy, 103, 776
ELISA, 266, 628
embryogenic callus, 266
Endothia gyrosa, 211
epidemiology, 234, 317, 433, 725
epoxiconazole, 190
ergot, 681
Erysiphe cichoracearum, 690
Erysiphe graminis, 317, 552
Eucalyptus smithii, 97
Eucalyptus, 587
evolution, 732
exponential growth rate, 363
eyespot, 356

F_1 hybrid, 97
Fabaceae, 433
FAMEs, 653
Ficus spp., 620
fluquinconazole, 75
fungal pathogen, 690
fungi, 593
fungicicde, 244
fungicidal effects, 190
fungicide resistance, 302, 601
fungicide, 75, 453
Furovirus, 761
fusarium crown rot, 371
fusarium dry rot, 489
Fusarium solani, 258

gas chromatography - flame ionization detection, 489
gas chromatography - mass spectrometry, 489
genetic diversity, 690
genotype-environment interaction, 281
genotypes, 747
geostatistics, 53
germplasm, 681
GLC, 653
glucose-6-phosphate isomerase, 302
grapevine, 427
growth rate, 497

Helminthosporium solani, 141
Heterobasidion annosum, 230
Heterodera avenae complex, 270
heterokaryon, 371
honeydew, 681
host specificity, 433
hydroponics, 503, 702

immigration, 317
in plata detection, 609
incubation period, 28
induced disease resistance, 230
induced systematic resistance, 579
infection conditions, 28
infectious period, 154
infectivity, 420
ink disease, 90
inoculations, 111
integrated pest management, 579
Ipomovirus, 527, 768
isoelectrofocusing, 270
isolation *in vitro*, 420
isomarticin, 258
isozymes, 270

jarrah forest, 587

Karnal bunt, 470

late blight, 489
latent period, 42, 154, 363
Leptographium, 111
Leptosphaeria maculans, 28
lipid peroxidation, 593
Loganberry, 747
luteovirus, 676
Lycopersicon esculentum, 477

marker-assisted selection, 637
mating type, 90, 427
median tetrad analysis, 325
metabolic profiling, 537
metalaxyl, 302
molecular markers, 165
Monilinia fructigena, 561, 569
mono/ polycyclic process, 53
mortality, 561
mustard, 69
Mycosphaerella graminicola, 325, 339
Myrtaceae, 725

naphthazarins, 258
Nectria fuckeliana, 230
nematicides, 509
Neovossia indica, 470
nit mutants, 371
Norway spruce, 230, 497
nutrient-film technique, 503

Oidium anacardii, 234
oilseed rape, 69
Olea europaea, 609
olive, 609
onions, 249
Oomycota, 433
oospores, 427
Ophiostoma, 111

opines, 620
oxidative stress, 593
oxygen deficiency, 497

P. *citricola*, 90
P. *gonapodyides*, 90
Pahseolus vulgaris, 443
partial resistance, 546
passage cells, 249
pathogenicity, 90, 477, 497
pathotype, 174
pathotypes, 181, 609
PCR detection, 782
PCR identification, 420
PCR, 124, 515, 609, 628
PCR-RFLPs, 747
pea common root rot, 433
pear, 569
pearl millet, 310
peiodical regression, 725
peroxidase phenols, 470
pH, 503
Phaeoisariopsis griseola, 443
phage bath, 666
Phaseolus vulgaris, 103
phoma leaf spot, 10, 28
Phoma lingam, 10, 69
phosphate, 83
phosphite, 83, 587
photosynthesis, 443
Phytophthora cactorum, 90
Phytophthora cinnamoni, 97, 587
Phytophthora infestans, 281, 292
phytophthora root rot, 702
Phytophthora, 83, 218
phytoplasma, 776
phytoplasmas, 130
pink rot, 302
Polerovirus, 628
polycyclic disease, 42
polymerase chain reaction, 130, 776
population diversity, 206
population genetics, 317
postinfection period, 154
potato early dying syndrome, 483
potato late blight, 281
potato store disease, 141
potato tuber blight, 292
potato tuber, 302, 666
potato virus Y, 124
Potyviridae, 527
powdery mildew, 234, 244, 552
Pratylenchus penetrans, 509
predawn leaf water potential, 708
primary inoculum, 317
proline, 702
Pseudocercosporella herpotrichoides, 356
Puccinia lagenphorae, 363
Puccinia psidii, 725
Puccinia triticina, 546
Pyrenophora graminea, 347

QTL analysis, 690
quantitative field resistance, 281
quarantine, 628
Quercus ilex, 708
Quercus rubra, 708
Quercus suber, 708
Quercus, 218

race-specific resistance, 69
RAPD, 97, 165, 174
RAPD-PCR analysis, 117
rapid PCR detection ,347
rasberry, 747
Rehmannia glutinosa, 776
REP-PCR, 397
resistance, 234, 637, 681
restriction fragment length polymorphism, 130, 776
RFLP, 211

R-gene pyramiding, 690
Rhizoctonia cerealis, 463
Rhizoctonia solani, 53, 593
ribosomal DNA sequence, 211
rice sheath blight, 53
root fungi, 75
root-knot nematode, 637
rootstocks, 258
RT-PCR detection, 768
RT-PCR, 628, 761

Saccharum spp. hybrids, 676
sampling, 755
saprophytic effects, 190
SCAR, 347
Sclerospora graminicola, 310
sclerotia, 593
Sclerotium cepivorum, 249
Scots pine, 111
SDS-PAGE, 653
seasonal variation, 218
seed germination, 117
seed health test, 347
seed treatment, 75
Senecio vulgaris, 363
septoria tritici blotch, 325, 339
serology, 266
Serology, 653
sexual recombination, 97
shearing forces, 310
silicon, 537
silver scurf, 141
single-spore isolates, 165
Sirex woodwasp, 206
slow-rusting resistance, 546
soft rot, 117
soil amendments, 537
soil-borne disease, 433
soilborne inoculum, 53
Soil-borne wheat mosaic virus, 761
Solanum phureja, 281
Solanum tuberosum, 477, 483, 489
somatic embryos, 266
sorghum, 681
spatiotempral analysis, 53
splash dispersal, 42
Spongospora subterranea f.sp. *subterranea*, 420
sporangia, 310, 587
spore germination, 103, 601
spore production, 363
sporulation, 561
spray-inoculation, 310
stability of disease resistance, 292
stem canker, 28, 69
stem-based disease, 463
Sternorryncha, 782
stigma, 266
stomatal conductance, 708
stubble, 356
style, 266
sugarcane yellow leaf syndrome, 676
Sugarcane yellow leaf virus, 628, 676
sugarcane yellows phytoplasma, 628
surveys, 453
susceptibility, 154
sustainable agriculture, 509
Syzygium jambos, 725

take-all, 75
Tapesia acuformis, 356
Tapesia spp., 463
Tapesia yallundae, 356
taxonomy, 270
Tayberry, 747
TBZ resistance, 141
thiabendazole, 141
Tilletia controversa, 755
tissue culture, 483, 676
tomato bacterial canker, 503
tomato, 371
tomato, 702
Tomicus, 111
transfer function, 717
Trichoderma koningii, 249
Triticum aestivum, 470, 537, 546
Triticum durum, 470
tropical fungi, 725
TSWV, 402
tuber infection, 402
tuber yields, 509

urediniospores, 181
Uromyces appendiculatus, 443

variability, 755
variation ,174
VCG, 477
vegetative compatability, 371
vegetative incompatibility, 206
velocity of focus expansion, 363
Verticillium dahliae, 609
verticillium wilt, 477, 483
virulence, 165, 732
virus distribution, 747
virus transmission, 732
virus transport, 402
visual disease assessment, 463
Vitis, 427, 515

water relations, 708
wheat stripe rust, 174
wheat yellow rust, 174, 181
wheat, 75, 325, 339, 470, 755
white rot, 249
winter oilseed rape, 28
winter wheat, 190, 453
wounding, 569

yield loss, 443
yield response, 244
yield, 190
Yr genes, 181

zoospores, 310, 587